

A MATRIX APPROACH TO THE RATIONAL INVARIANTS OF CERTAIN CLASSICAL GROUPS OVER FINITE FIELDS OF CHARACTERISTIC TWO

ZHONGMING TANG AND ZHE-XIAN WAN

ABSTRACT. Let $\mathbb{F}_q$ be a finite field of characteristic two and $\mathbb{F}_q(X_1, \dots, X_n)$ a rational function field. We use matrix methods to obtain explicit transcendental bases of the invariant subfields of orthogonal groups and pseudo-symplectic groups on $\mathbb{F}_q(X_1, \dots, X_n)$ over $\mathbb{F}_q$.

1. INTRODUCTION

Let $\mathbb{F}_q$ be a finite field, $n \geq 1$ an integer and $\mathrm{GL}_n(\mathbb{F}_q)$ the general linear group. For any $T \in \mathrm{GL}_n(\mathbb{F}_q)$, T induces an $\mathbb{F}_q$ -linear action σ_T on the rational function field $\mathbb{F}_q(X_1, \dots, X_n)$ defined by $\sigma_T(f(X_1, \dots, X_n)) = f(\sigma_T(X_1), \dots, \sigma_T(X_n))$ for all $f(X_1, \dots, X_n) \in \mathbb{F}_q(X_1, \dots, X_n)$, where

$$(\sigma_T(X_1), \dots, \sigma_T(X_n)) = (X_1, \dots, X_n)T.$$

Let G be a subgroup of $\mathrm{GL}_n(\mathbb{F}_q)$. The invariant subfield of G on $\mathbb{F}_q(X_1, \dots, X_n)$ is

$$\mathbb{F}_q(X_1, \dots, X_n)^G = \{f \in \mathbb{F}_q(X_1, \dots, X_n) : \sigma_T(f) = f \text{ for all } T \in G\}.$$

Then $\mathbb{F}_q(X_1, \dots, X_n)$ is a finite Galois extension of $\mathbb{F}_q(X_1, \dots, X_n)^G$. One asks when $\mathbb{F}_q(X_1, \dots, X_n)^G$ is purely transcendental over $\mathbb{F}_q$ for a classical group G .

When $G = \mathrm{GL}_n(\mathbb{F}_q)$, Dickson[6] gave an affirmative answer:

$$\mathbb{F}_q(X_1, \dots, X_n)^{\mathrm{GL}_n(\mathbb{F}_q)} = \mathbb{F}_q(C_{n0}, C_{n1}, \dots, C_{n,n-1}),$$

where $C_{ni} = \frac{D_{ni}}{D_{nn}}$, $i = 0, 1, \dots, n-1$, and

$$D_{ni} = \det \begin{pmatrix} X_1 & X_1^q & \dots & \widehat{X_1^{q^i}} & \dots & X_1^{q^n} \\ X_2 & X_2^q & \dots & \widehat{X_2^{q^i}} & \dots & X_2^{q^n} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ X_n & X_n^q & \dots & \widehat{X_n^{q^i}} & \dots & X_n^{q^n} \end{pmatrix}, \quad i = 0, 1, \dots, n.$$

Relative recently, Chu[3] considered rational invariants of orthogonal groups and obtained similar results for $n = 2, 3$. Afterwards, Cohen[5] showed the result is true when $n = 4$, and Carlisle and Kropholler[2] solved the general case. But they all assumed that the characteristic of $\mathbb{F}_q$ is odd. The case of characteristic two was settled by Rajaei[7] using quadratic form language.

Key words and phrases. rational invariant, rational function field, orthogonal group, pseudo-symplectic group.

2000 Mathematics Subject Classification. 20G40, 13A50, 16W22.

Both authors are supported by the National Natural Science Foundation of China.

Suppose that the characteristic of $\mathbb{F}_q$ is two. In this paper, we will follow the route of Carlisle and Kropholler[2] and Chu[4] to use matrix methods to give explicit transcendental bases of the invariant subfields of orthogonal groups on $\mathbb{F}_q(X_1, \dots, X_n)$ over $\mathbb{F}_q$. Our transcendental bases are different from that of [7] in some cases. Moreover, we also obtain explicit transcendental bases of the invariant subfields of pseudo-symplectic groups on $\mathbb{F}_q(X_1, \dots, X_n)$ over $\mathbb{F}_q$.

Following Chu[4], some results in [2] and [4] can be described as follows. Let $A \in \text{GL}_n(\mathbb{F}_q)$ be a symmetric matrix and

$$G_A = \{T \in \text{GL}_n(\mathbb{F}_q) : TA^tT = A\},$$

where tT denotes the transpose of T . Define

$$P_{nk} = (X_1, \dots, X_n)A \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_n^{q^k} \end{pmatrix}, \quad k = 0, 1, 2, \dots$$

Then

$$P_{nk} \in \mathbb{F}_q(X_1, \dots, X_n)^{G_A}, \quad k = 0, 1, 2, \dots,$$

and

$$\mathbb{F}_q(X_1, \dots, X_n)^{G_A} = \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{nn}),$$

(cf., the proof of [4, Theorem]). Here there is no restriction on the characteristic of $\mathbb{F}_q$. When the characteristic of $\mathbb{F}_q$ is odd,

$$\mathbb{F}_q(X_1, \dots, X_n)^{G_A} = \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{n,n-1}),$$

i.e., $P_{nn} \in \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{n,n-1})$. The restriction on the characteristic of $\mathbb{F}_q$ is crucial in their proof. Similar problems for rational invariants of unitary groups and symplectic groups have been also solved (cf., [2] and [4]), where there is no restriction on the characteristic of $\mathbb{F}_q$. For example, for the symplectic group $Sp_{2\nu}(\mathbb{F}_q)$ of degree 2ν over $\mathbb{F}_q$ of any characteristic, we have

$$\mathbb{F}_q(X_1, \dots, X_n)^{Sp_{2\nu}(\mathbb{F}_q)} = \mathbb{F}_q(P_{n1}, P_{n2}, \dots, P_{nn}).$$

Let $A = (a_{ij})$ be an $n \times n$ matrix over $\mathbb{F}_q$. A is said to be alternate if $a_{ij} = -a_{ji}$ and $a_{ii} = 0$. Let A and B be two $n \times n$ matrices, we write $A \equiv B$ if $A - B$ is alternate. The identity matrix of rank n will be denoted by $I^{(n)}$.

Suppose that the characteristic of $\mathbb{F}_q$ is two. Then, up to isomorphisms, orthogonal groups over $\mathbb{F}_q$ are just the following three types:

$$O_n(\mathbb{F}_q, G) = \{T \in \text{GL}_n(\mathbb{F}_q) : TG^tT \equiv G\},$$

where G is one of the following

$$\begin{pmatrix} 0 & I^{(\nu)} \\ & 0 \end{pmatrix}; \begin{pmatrix} 0 & I^{(\nu)} \\ & 0 & 1 \end{pmatrix}; \begin{pmatrix} 0 & I^{(\nu-1)} & & \\ & 0 & & \\ & & \alpha & 1 \\ & & & \alpha \end{pmatrix},$$

with $\alpha \in \mathbb{F}_q \setminus \{x + x^2 : x \in \mathbb{F}_q\}$, which will be denoted by $G_{2\nu}$, $G_{2\nu+1}$, $G_{2(\nu-1)+2}$, respectively. While the pseudo-symplectic group with respect to non-alternate symmetric matrices are just the following two types:

$$Ps_n(\mathbb{F}_q, S) = \{T \in \text{GL}_n(\mathbb{F}_q) : TS^tT = S\},$$

where S is one of the following

$$\begin{pmatrix} 0 & I^{(\nu)} & & \\ I^{(\nu)} & 0 & & \\ & & 1 & \\ & & & 1 \end{pmatrix}; \begin{pmatrix} 0 & I^{(\nu)} & & \\ I^{(\nu)} & 0 & & \\ & & 0 & 1 \\ & & 1 & 1 \end{pmatrix},$$

which will be denoted by $S_{2\nu+1}$, $S_{2\nu+2}$, respectively, cf., Wan[8].

For the rational invariants of orthogonal groups $O_n(\mathbb{F}_q, G)$ where G is $G_{2\nu}$ or $G_{2(\nu-1)+2}$, let $A = G + {}^tG$ and G_A , P_{nk} for $k > 0$ be defined as before but define

$$P_{n0} = (X_1, \dots, X_n)G \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix},$$

(note that the former definition gives $P_{n0} = 0$). Then $O_n(\mathbb{F}_q, G)$ is a subgroup of G_A , which is isomorphic to $Sp_{2\nu}(\mathbb{F}_q)$, and

$$\mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)} = \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{nn}).$$

The main part of the paper is devoted to show that $P_{nn} \in \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{n,n-1})$. We need to find a polynomial identity like $aP_{nn} = b$ where $a, b \in \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{n,n-1})$ but $a \neq 0$ as done in odd characteristic case in [2] and [4]. However, such an identity in odd characteristic case becomes trivial in characteristic two ($a = 0!$). To overcome this difficulty, we first go to characteristic zero, get some identity and cancel the factor 2. Then, returning to characteristic two, we obtain a required identity. In order to do this, we need some properties of determinants which form section 2. The results of rational invariants of another orthogonal group and pseudo-symplectic groups are obtained similarly. Section 3 consists of results about orthogonal groups and section 4 discusses pseudo-symplectic groups.

2. PRELIMINARIES

The following properties of determinants are useful in our discussion. Although some of them are probably known, for the reader's convenience, we give their proofs. The entries of matrices will be considered as indeterminates.

Lemma 2.1. *If n is odd, then, in $\mathbb{Z}[a_{ij} : 1 \leq i < j \leq n]$,*

$$\det \begin{pmatrix} 0 & a_{12} & \dots & a_{1n} \\ a_{12} & 0 & \dots & a_{2n} \\ & & \dots & \dots \\ a_{1n} & a_{2n} & \dots & 0 \end{pmatrix} \equiv 0 \pmod{2}.$$

Proof. It is because that, in $\mathbb{Z}_2[a_{ij} : 1 \leq i < j \leq n]$,

$$\det \begin{pmatrix} 0 & a_{12} & \dots & a_{1n} \\ a_{12} & 0 & \dots & a_{2n} \\ & & \dots & \dots \\ a_{1n} & a_{2n} & \dots & 0 \end{pmatrix} = 0.$$

□

Lemma 2.2. *If n is even, then, in $\mathbb{Z}[a_{ij} : 1 \leq i \leq j \leq n]$,*

$$\det \begin{pmatrix} 2a_{11} & a_{12} & \dots & a_{1n} \\ a_{12} & 2a_{22} & \dots & a_{2n} \\ & \dots & \dots & \\ a_{1n} & a_{2n} & \dots & 2a_{nn} \end{pmatrix} \equiv \det \begin{pmatrix} 0 & a_{12} & \dots & a_{1n} \\ a_{12} & 0 & \dots & a_{2n} \\ & \dots & \dots & \\ a_{1n} & a_{2n} & \dots & 0 \end{pmatrix} \pmod{4}.$$

Proof. Since

$$\begin{aligned} & \det \begin{pmatrix} 2a_{11} & a_{12} & \dots & a_{1n} \\ a_{12} & 2a_{22} & \dots & a_{2n} \\ & \dots & \dots & \\ a_{1n} & a_{2n} & \dots & 2a_{nn} \end{pmatrix} \\ &= 2a_{11} \det \begin{pmatrix} 2a_{22} & a_{23} & \dots & a_{2n} \\ a_{23} & 2a_{33} & \dots & a_{3n} \\ & \dots & \dots & \\ a_{2n} & a_{3n} & \dots & 2a_{nn} \end{pmatrix} + \det \begin{pmatrix} 0 & a_{12} & \dots & a_{1n} \\ a_{12} & 2a_{22} & \dots & a_{2n} \\ & \dots & \dots & \\ a_{1n} & a_{2n} & \dots & 2a_{nn} \end{pmatrix} \\ &\equiv 2a_{11} \det \begin{pmatrix} 0 & a_{23} & \dots & a_{2n} \\ a_{23} & 0 & \dots & a_{3n} \\ & \dots & \dots & \\ a_{2n} & a_{3n} & \dots & 0 \end{pmatrix} + \det \begin{pmatrix} 0 & a_{12} & \dots & a_{1n} \\ a_{12} & 2a_{22} & \dots & a_{2n} \\ & \dots & \dots & \\ a_{1n} & a_{2n} & \dots & 2a_{nn} \end{pmatrix} \pmod{4}, \end{aligned}$$

it follows from 2.1 that

$$\det \begin{pmatrix} 2a_{11} & a_{12} & \dots & a_{1n} \\ a_{12} & 2a_{22} & \dots & a_{2n} \\ & \dots & \dots & \\ a_{1n} & a_{2n} & \dots & 2a_{nn} \end{pmatrix} \equiv \det \begin{pmatrix} 0 & a_{12} & \dots & a_{1n} \\ a_{12} & 2a_{22} & \dots & a_{2n} \\ & \dots & \dots & \\ a_{1n} & a_{2n} & \dots & 2a_{nn} \end{pmatrix} \pmod{4}.$$

Then the result follows by induction. $\square$

Lemma 2.3. *If $n = 2\nu$ is even, then, in $\mathbb{Z}[a_{ij} : 1 \leq i < j \leq n]$,*

$$\det \begin{pmatrix} 0 & a_{12} & a_{13} & \dots & a_{1n} \\ -a_{12} & 0 & a_{23} & \dots & a_{2n} \\ -a_{13} & -a_{23} & 0 & \dots & a_{3n} \\ \dots & \dots & \dots & \dots & \\ -a_{1n} & -a_{2n} & -a_{3n} & \dots & 0 \end{pmatrix} \equiv (-1)^\nu \det \begin{pmatrix} 0 & a_{12} & a_{13} & \dots & a_{1n} \\ a_{12} & 0 & a_{23} & \dots & a_{2n} \\ a_{13} & a_{23} & 0 & \dots & a_{3n} \\ \dots & \dots & \dots & \dots & \\ a_{1n} & a_{2n} & a_{3n} & \dots & 0 \end{pmatrix} \pmod{4}.$$

Proof. By induction on ν . The case $\nu = 1$ is clear. Suppose that $\nu > 1$ and the result is true for $\nu - 1$. Set

$$A = \begin{pmatrix} 0 & a_{12} & a_{13} & \dots & a_{1n} \\ -a_{12} & 0 & a_{23} & \dots & a_{2n} \\ -a_{13} & -a_{23} & 0 & \dots & a_{3n} \\ \dots & \dots & \dots & \dots & \\ -a_{1n} & -a_{2n} & -a_{3n} & \dots & 0 \end{pmatrix}, B = \begin{pmatrix} 0 & a_{12} & a_{13} & \dots & a_{1n} \\ a_{12} & 0 & a_{23} & \dots & a_{2n} \\ a_{13} & a_{23} & 0 & \dots & a_{3n} \\ \dots & \dots & \dots & \dots & \\ a_{1n} & a_{2n} & a_{3n} & \dots & 0 \end{pmatrix}.$$

Let us consider a_{ij} as indeterminates and work in the field $\mathbb{Q}(a_{ij} : 1 \leq i < j \leq n)$. Let E be the matrix obtained from the $2\nu \times 2\nu$ identity matrix by replacing the zeros at

$(2, 3), (2, 4), \dots, (2, n)$ positions by $-\frac{a_{13}}{a_{12}}, -\frac{a_{14}}{a_{12}}, \dots, -\frac{a_{1n}}{a_{12}}$, respectively. Then

$${}^tEAE = \begin{pmatrix} 0 & a_{12} & 0 & \dots & 0 \\ -a_{12} & 0 & a_{23} & \dots & a_{2n} \\ 0 & -a_{23} & & & \\ \dots & & A_1 & & \\ 0 & -a_{2n} & & & \end{pmatrix}, {}^tEBE = \begin{pmatrix} 0 & a_{12} & 0 & \dots & 0 \\ a_{12} & 0 & a_{23} & \dots & a_{2n} \\ 0 & a_{23} & & & \\ \dots & & B_1 & & \\ 0 & a_{2n} & & & \end{pmatrix},$$

where

$$A_1 = \begin{pmatrix} 0 & a'_{34} & \dots & a'_{3n} \\ -a'_{34} & 0 & \dots & a'_{4n} \\ \dots & \dots & \dots & \dots \\ -a'_{3n} & -a'_{4n} & \dots & 0 \end{pmatrix}, B_1 = \begin{pmatrix} a''_{33} & a''_{34} & \dots & a''_{3n} \\ a''_{34} & a''_{44} & \dots & a''_{4n} \\ \dots & \dots & \dots & \dots \\ a''_{3n} & a''_{4n} & \dots & a''_{nn} \end{pmatrix},$$

and $a'_{ij} = a_{ij} - \frac{a_{1i}}{a_{12}}a_{2j} + \frac{a_{1j}}{a_{12}}a_{2i}$, $a''_{ij} = -2\frac{a_{1j}}{a_{12}}a_{2i} + a'_{ij}$ for $i < j$, $a''_{ii} = -2\frac{a_{1i}}{a_{12}}a_{2i}$. Note that $a_{12}a'_{ij}, a_{12}a''_{ij} \in \mathbb{Z}[a_{ij} : 1 \leq i < j \leq n]$.

Clearly $\det(A) = a_{12}^2 \det(A_1)$ and $\det(B) = -a_{12}^2 \det(B_1)$. Let

$$B_2 = \begin{pmatrix} 0 & a'_{34} & \dots & a'_{3n} \\ a'_{34} & 0 & \dots & a'_{4n} \\ \dots & \dots & \dots & \dots \\ a'_{3n} & a'_{4n} & \dots & 0 \end{pmatrix}.$$

By induction assumption, $a_{12}^{n-2}(\det(A_1) - (-1)^{\nu-1} \det(B_2)) \in 4\mathbb{Z}[a_{ij} : 1 \leq i < j \leq n]$. Note that

$$\begin{aligned} & a_{12}^{n-4}(\det(A) - (-1)^\nu \det(B)) \\ &= a_{12}^{n-2}(\det(A_1) - (-1)^{\nu-1} \det(B_1)) \\ &= a_{12}^{n-2}(\det(A_1) - (-1)^{\nu-1} \det(B_2)) + (-1)^{\nu-1} a_{12}^{n-2}(\det(B_2) - \det(B_1)). \end{aligned}$$

Thus it is enough to show that $a_{12}^{n-2}(\det(B_1) - \det(B_2)) \in 4\mathbb{Z}[a_{ij} : 1 \leq i < j \leq n]$. Let $c_{ij} = a'_{ij}$ for $3 \leq i < j \leq n$, and $d_{ij} = -\frac{a_{1i}}{a_{12}}a_{2j}$ for $3 \leq i \leq j \leq n$. Then

$$\begin{aligned} B_1 &= \begin{pmatrix} 2d_{33} & 2d_{34} + c_{34} & \dots & 2d_{3n} + c_{3n} \\ 2d_{34} + c_{34} & 2d_{44} & \dots & 2d_{4n} + c_{4n} \\ & \dots & \dots & \\ 2d_{3n} + c_{3n} & 2d_{4n} + c_{4n} & \dots & 2d_{nn} \end{pmatrix}, \\ B_2 &= \begin{pmatrix} 0 & c_{34} & \dots & c_{3n} \\ c_{34} & 0 & \dots & c_{4n} \\ \dots & \dots & \dots & \dots \\ c_{3n} & c_{4n} & \dots & 0 \end{pmatrix}. \end{aligned}$$

Let us show that $\det(B_1) - \det(B_2) \in 4\mathbb{Z}[d_{ij}, c_{ij} : 3 \leq i \leq j \leq n]$, where $c_{ii} = 0$, then the lemma follows.

Note that, by 2.2,

$$\det(B_1) - \det \begin{pmatrix} 0 & 2d_{34} + c_{34} & \dots & 2d_{3n} + c_{3n} \\ 2d_{34} + c_{34} & 0 & \dots & 2d_{4n} + c_{4n} \\ & \dots & \dots & \\ 2d_{3n} + c_{3n} & 2d_{4n} + c_{4n} & \dots & 0 \end{pmatrix} \in 4\mathbb{Z}[d_{ij}, c_{ij} : 3 \leq i \leq j \leq n],$$

hence, we may assume that $d_{ii} = 0$, $i = 3, \dots, n$. Then, in $\mathbb{Z}[d_{ij}, c_{ij} : 3 \leq i \leq j \leq n]$,

$$\begin{aligned}
& \det(B_1) \\
&= \det \begin{pmatrix} 0 & 2d_{34} & \dots & 2d_{3n} \\ 2d_{34} & 0 & \dots & 2d_{4n} + c_{4n} \\ & \dots & \dots & \\ 2d_{3n} & 2d_{4n} + c_{4n} & \dots & 0 \end{pmatrix} + \det \begin{pmatrix} 0 & 2d_{34} & \dots & 2d_{3n} \\ c_{34} & 0 & \dots & 2d_{4n} + c_{4n} \\ & \dots & \dots & \\ c_{3n} & 2d_{4n} + c_{4n} & \dots & 0 \end{pmatrix} \\
&+ \det \begin{pmatrix} 0 & c_{34} & \dots & c_{3n} \\ 2d_{34} & 0 & \dots & 2d_{4n} + c_{4n} \\ & \dots & \dots & \\ 2d_{3n} & 2d_{4n} + c_{4n} & \dots & 0 \end{pmatrix} + \det \begin{pmatrix} 0 & c_{34} & \dots & c_{3n} \\ c_{34} & 0 & \dots & 2d_{4n} + c_{4n} \\ & \dots & \dots & \\ c_{3n} & 2d_{4n} + c_{4n} & \dots & 0 \end{pmatrix} \\
&\equiv \det \begin{pmatrix} 0 & c_{34} & \dots & c_{3n} \\ c_{34} & 0 & \dots & 2d_{4n} + c_{4n} \\ & \dots & \dots & \\ c_{3n} & 2d_{4n} + c_{4n} & \dots & 0 \end{pmatrix} \pmod{4},
\end{aligned}$$

thus, continuing reductively we get that, in $\mathbb{Z}[d_{ij}, c_{ij} : 3 \leq i \leq j \leq n]$,

$$\det(B_1) \equiv \det \begin{pmatrix} 0 & c_{34} & \dots & c_{3n} \\ c_{34} & 0 & \dots & c_{4n} \\ & \dots & \dots & \\ c_{3n} & c_{4n} & \dots & 0 \end{pmatrix} \pmod{4},$$

as required. $\square$

Let $A = (a_{ij})$ be an $n \times n$ alternate matrix. When n is even, one can define the Pfaffian of A , denoted by $\text{Pf}(A)$, cf., [1, §5, no.2]. When n is odd, set $\text{Pf}(A) = 0$. The following properties of the Pfaffians are well-known.

- (i) $\det(A) = (\text{Pf}(A))^2$.
- (ii) Interchanging two rows and same columns simultaneously, the Pfaffian changes by -1 .
- (iii) Pfaffians can be developed along a row:

$$\text{Pf}(A) = \sum_{j=1}^n (-1)^{i+j-1} \sigma(i, j) a_{ij} \text{Pf}(A_{i,j}), \quad i = 1, \dots, n,$$

where $\sigma(i, j)$ is the sign of $j - i$ and $A_{i,j}$ is the matrix obtained from A by deleting its i -th and j -th rows and columns together.

We also need the following property of Pfaffians.

Lemma 2.4. *Let $n > 0$ be an even integer and C an $(n - 1) \times (n - 1)$ alternate matrix. Then, for any $a_{12}, \dots, a_{1n}, b_{12}, \dots, b_{1n}$,*

$$\text{Pf} \begin{pmatrix} 0 & a_{12} \cdots a_{1n} \\ -a_{12} & \\ \vdots & C \\ -a_{1n} & \end{pmatrix} \cdot \text{Pf} \begin{pmatrix} 0 & b_{12} \cdots b_{1n} \\ -b_{12} & \\ \vdots & C \\ -b_{1n} & \end{pmatrix} = \det \begin{pmatrix} 0 & a_{12} \cdots a_{1n} \\ -b_{12} & \\ \vdots & C \\ -b_{1n} & \end{pmatrix}.$$

Proof. Consider a_{1i} , b_{1j} and the non-diagonal elements of C as indeterminates and work in the rational field generated by them over $\mathbb{Q}$. Let

$$A = \begin{pmatrix} 0 & a_{12} & \cdots & a_{1n} \\ -a_{12} & & & \\ \vdots & & C & \\ -a_{1n} & & & \end{pmatrix}, B = \begin{pmatrix} 0 & b_{12} & \cdots & b_{1n} \\ -b_{12} & & & \\ \vdots & & C & \\ -b_{1n} & & & \end{pmatrix}, D = \begin{pmatrix} 0 & a_{12} & \cdots & a_{1n} \\ -b_{12} & & & \\ \vdots & & C & \\ -b_{1n} & & & \end{pmatrix}.$$

We first use induction on $\frac{n}{2}$ to show that $\det(A) \cdot \det(B) = (\det(D))^2$. When $n = 2$, it is clear. Now assume that $n \geq 4$. Write

$$C = \begin{pmatrix} C_{11} & C_{12} \\ -{}^t C_{12} & C_{22} \end{pmatrix},$$

where C_{11} is $(n-3) \times (n-3)$, C_{12} is $(n-3) \times 2$ and C_{22} is 2×2 , and

$$A_1 = (a_{12} \cdots a_{1,n-2}), A_2 = (a_{1,n-1} \ a_{1n}), B_1 = (b_{12} \cdots b_{1,n-2}), B_2 = (b_{1,n-1} \ b_{1n}).$$

Then

$$\begin{aligned} & \begin{pmatrix} 1 & & -A_2 C_{22}^{-1} \\ & I^{(n-3)} & -C_{12} C_{22}^{-1} \\ & & I^{(2)} \end{pmatrix} A \begin{pmatrix} 1 & & \\ & I^{(n-3)} & \\ C_{22}^{-1t} A_2 & C_{22}^{-1t} C_{12} & I^{(2)} \end{pmatrix} \\ &= \begin{pmatrix} 0 & A_1 + A_2 C_{22}^{-1t} C_{12} & 0 \\ -{}^t A_1 + C_{12} C_{22}^{-1t} A_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} & 0 \\ 0 & 0 & C_{22} \end{pmatrix}. \end{aligned}$$

Thus

$$\det(A) = \det(C_{22}) \cdot \det \begin{pmatrix} 0 & A_1 + A_2 C_{22}^{-1t} C_{12} \\ -{}^t A_1 + C_{12} C_{22}^{-1t} A_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} \end{pmatrix}.$$

Similarly,

$$\begin{aligned} \det(B) &= \det(C_{22}) \cdot \det \begin{pmatrix} 0 & B_1 + B_2 C_{22}^{-1t} C_{12} \\ -{}^t B_1 + C_{12} C_{22}^{-1t} B_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} \end{pmatrix} \\ \det(D) &= \det(C_{22}) \cdot \det \begin{pmatrix} A_2 C_{22}^{-1t} B_2 & A_1 + A_2 C_{22}^{-1t} C_{12} \\ -{}^t B_1 + C_{12} C_{22}^{-1t} B_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} \end{pmatrix}. \end{aligned}$$

Since $C_{11} + C_{12} C_{22}^{-1t} C_{12}$ is alternate of odd order $n-3$, its determinant is 0, hence

$$\begin{aligned} & \det \begin{pmatrix} A_2 C_{22}^{-1t} B_2 & A_1 + A_2 C_{22}^{-1t} C_{12} \\ -{}^t B_1 + C_{12} C_{22}^{-1t} B_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} \end{pmatrix} \\ &= \det \begin{pmatrix} 0 & A_1 + A_2 C_{22}^{-1t} C_{12} \\ -{}^t B_1 + C_{12} C_{22}^{-1t} B_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} \end{pmatrix}. \end{aligned}$$

By induction hypothesis

$$\begin{aligned} & \det \begin{pmatrix} 0 & A_1 + A_2 C_{22}^{-1t} C_{12} \\ -{}^t A_1 + C_{12} C_{22}^{-1t} A_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} \end{pmatrix} \cdot \det \begin{pmatrix} 0 & B_1 + B_2 C_{22}^{-1t} C_{12} \\ -{}^t B_1 + C_{12} C_{22}^{-1t} B_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} \end{pmatrix} \\ &= \left(\det \begin{pmatrix} 0 & A_1 + A_2 C_{22}^{-1t} C_{12} \\ -{}^t B_1 + C_{12} C_{22}^{-1t} B_2 & C_{11} + C_{12} C_{22}^{-1t} C_{12} \end{pmatrix} \right)^2. \end{aligned}$$

Therefore $\det(A) \cdot \det(B) = (\det(D))^2$. It follows that $\text{Pf}(A) \cdot \text{Pf}(B) = \pm \det(D)$.

Now, we use one term to determine the sign in front of $(\det(D))^2$. Let $C = (c_{ij})$. Since the sign of term $a_{12}c_{23}c_{45} \cdots c_{n-2,n-1}$ in $\text{Pf}(A)$ is 1, the sign of term $b_{12}c_{23}c_{45} \cdots c_{n-2,n-1}$ in $\text{Pf}(B)$ is 1 and the sign of the corresponding term $a_{12}b_{12}c_{23}^2c_{45}^2 \cdots c_{n-2,n-1}^2$ in $\det(D)$ is also 1, it follows that the above sign in front of $\det(D)$ is 1, i.e., $\text{Pf}(A) \cdot \text{Pf}(B) = \det(D)$. $\square$

3. RATIONAL INVARIANTS OF ORTHOGONAL GROUPS

Consider first the case when $n = 2\nu$ is even. We begin by deducing some polynomial identities over $\mathbb{Z}$. Let q be a power of a prime p and $f = \sum a_{\alpha_1 \dots \alpha_n} X_1^{\alpha_1} \cdots X_n^{\alpha_n} \in \mathbb{Z}[X_1, \dots, X_n]$. Denote $\sum a_{\alpha_1 \dots \alpha_n} X_1^{\alpha_1 q^r} \cdots X_n^{\alpha_n q^r}$ by $f^{(q^r)}$. For any matrix $C = (c_{ij})$ over $\mathbb{Z}[X_1, \dots, X_n]$, let $C^{(q^r)} = (c_{ij}^{(q^r)})$. Embed $\mathbb{Z}_p$ in $\mathbb{F}_q$ as a subring. Under the natural map from $\mathbb{Z}$ to $\mathbb{Z}_p$, we will use the same f to denote the natural image of $f \in \mathbb{Z}[X_1, \dots, X_n]$ in $\mathbb{Z}_p[X_1, \dots, X_n]$. By our convention, $f^{(q^r)} = f^{q^r}$ in $\mathbb{Z}_p[X_1, \dots, X_n]$ for any $f \in \mathbb{Z}[X_1, \dots, X_n]$. Let $G = G_{2\nu}$ or $G_{2(\nu-1)+2}$, but now the α in $G_{2(\nu-1)+2}$ is regarded as an indeterminate. Set $\alpha = 0$ when $G = G_{2\nu}$. Let $A = G + {}^t G$ and $\bar{A} = G - {}^t G$. Then $\det(A) = (-1)^{\nu-1} 4\alpha^2 + (-1)^\nu$ and $\det(\bar{A}) = 1$ thus both A and $\bar{A}$ are nonsingular in $\mathbb{Z}(\alpha)$ and also in $\mathbb{Z}_p(\alpha)$. In $\mathbb{Z}[X_1, \dots, X_n]$, define

$$P_{n0} = (X_1, \dots, X_n)G \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix}, \quad (1)$$

$$P_{nk} = (X_1, \dots, X_n)A \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_n^{q^k} \end{pmatrix}, \quad k = 1, 2, \dots \quad (2)$$

Let

$$B = \begin{pmatrix} X_1 & X_2 & \cdots & X_n \\ X_1^q & X_2^q & \cdots & X_n^q \\ \vdots & \vdots & \ddots & \vdots \\ X_1^{q^{n-1}} & X_2^{q^{n-1}} & \cdots & X_n^{q^{n-1}} \end{pmatrix}.$$

and

$$P = \begin{pmatrix} 0 & P_{n1} & P_{n2} & \cdots & P_{n,n-1} \\ -P_{n1} & 0 & P_{n1}^{(q)} & \cdots & P_{n,n-2}^{(q)} \\ -P_{n2} & -P_{n1}^{(q)} & 0 & \cdots & P_{n,n-3}^{(q^2)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -P_{n,n-1} & -P_{n,n-2}^{(q)} & -P_{n,n-3}^{(q^2)} & \cdots & 0 \end{pmatrix}.$$

Then we have

Lemma 3.1.

$$\text{Pf}(P) \cdot \text{Pf}(P^{(q)}) = \det \begin{pmatrix} 0 & P_{n1} & P_{n2} & \cdots & P_{n,n-1} \\ -P_{n,n-1}^{(q)} & 0 & P_{n1}^{(q)} & \cdots & P_{n,n-2}^{(q)} \\ -P_{n,n-2}^{(q^2)} & -P_{n1}^{(q)} & 0 & \cdots & P_{n,n-3}^{(q^2)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -P_{n1}^{(q^{n-1})} & -P_{n,n-2}^{(q)} & -P_{n,n-3}^{(q^2)} & \cdots & 0 \end{pmatrix}.$$

Proof. Note that

$$\begin{aligned}
\text{Pf}(P^{(q)}) &= \text{Pf} \begin{pmatrix} 0 & P_{n1}^{(q)} & P_{n2}^{(q)} & \cdots & P_{n,n-1}^{(q)} \\ -P_{n1}^{(q)} & 0 & P_{n1}^{(q^2)} & \cdots & P_{n,n-2}^{(q^2)} \\ -P_{n2}^{(q)} & -P_{n1}^{(q^2)} & 0 & \cdots & P_{n,n-3}^{(q^3)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -P_{n,n-1}^{(q)} & -P_{n,n-2}^{(q^2)} & -P_{n,n-3}^{(q^3)} & \cdots & 0 \end{pmatrix} \\
&= -\text{Pf} \begin{pmatrix} 0 & -P_{n,n-1}^{(q)} & -P_{n,n-2}^{(q^2)} & \cdots & -P_{n1}^{(q^{n-1})} \\ P_{n,n-1}^{(q)} & 0 & P_{n1}^{(q)} & \cdots & P_{n,n-2}^{(q)} \\ P_{n,n-2}^{(q^2)} & -P_{n1}^{(q)} & 0 & \cdots & P_{n,n-3}^{(q^2)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ P_{n1}^{(q^{n-1})} & -P_{n,n-2}^{(q)} & -P_{n,n-3}^{(q^2)} & \cdots & 0 \end{pmatrix},
\end{aligned}$$

by property (ii) of Pfaffians. Then, our lemma follows from 2.4. $\square$

Let $E = (e_{ij})$ be an $m \times m$ matrix over $\mathbb{Z}[X_1, \dots, X_n]$. For convenience, we will use $\mathbb{Z}[E]$ to denote $\mathbb{Z}[e_{ij} : i, j = 1, \dots, m]$.

Lemma 3.2. *There is a polynomial $S \in \mathbb{Z}[X_1, \dots, X_n]$ such that*

$$\det(B) \equiv 2S + \text{Pf}(P) \pmod{4}$$

and

$$S(S + \text{Pf}(P)) \equiv f \pmod{2}$$

for some $f \in \mathbb{Z}[\alpha, P_{n0}, P_{n0}^{(q)}, \dots, P_{n0}^{(q^{n-1})}, P]$.

Proof. Let

$$\begin{aligned}
\overline{P}_{nk} &= (X_1, \dots, X_n) \overline{A} \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_n^{q^k} \end{pmatrix}, \\
P'_{nk} &= (X_1, \dots, X_n)^t G \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_n^{q^k} \end{pmatrix}, \quad k = 1, 2, \dots,
\end{aligned}$$

and

$$\overline{P} = \begin{pmatrix} 0 & \overline{P}_{n1} & \overline{P}_{n2} & \cdots & \overline{P}_{n,n-1} \\ -\overline{P}_{n1} & 0 & \overline{P}_{n1}^{(q)} & \cdots & \overline{P}_{n,n-2}^{(q)} \\ -\overline{P}_{n2} & -\overline{P}_{n1}^{(q)} & 0 & \cdots & \overline{P}_{n,n-3}^{(q^2)} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ -\overline{P}_{n,n-1} & -\overline{P}_{n,n-2}^{(q)} & -\overline{P}_{n,n-3}^{(q^2)} & \cdots & 0 \end{pmatrix}.$$

Clearly, $B\bar{A}^t B = \bar{P}$. Since $\det(\bar{A}) = 1$, $(\det(B))^2 = \det(B\bar{A}^t B) = \det(\bar{P}) = (\text{Pf}(\bar{P}))^2$. Thus $\det(B) = \pm \text{Pf}(\bar{P})$. Clearly, $\bar{P}_{nk} = P_{nk} - 2P'_{nk}$. Therefore

$$\begin{aligned} & \text{Pf}(\bar{P}) \\ = & \text{Pf} \begin{pmatrix} 0 & P_{n1} - 2P'_{n1} & P_{n2} - 2P'_{n2} & \cdots & P_{n,n-1} - 2P'_{n,n-1} \\ -P_{n1} + 2P'_{n1} & 0 & P_{n1}^{(q)} - 2P_{n1}'^{(q)} & \cdots & P_{n,n-2}^{(q)} - 2P_{n,n-2}'^{(q)} \\ -P_{n2} + 2P'_{n2} & -P_{n1}^{(q)} + 2P_{n1}'^{(q)} & 0 & \cdots & P_{n,n-3}^{(q^2)} - 2P_{n,n-3}'^{(q^2)} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ -P_{n,n-1} + 2P'_{n,n-1} & -P_{n,n-2}^{(q)} + 2P_{n,n-2}'^{(q)} & -P_{n,n-3}^{(q^2)} + 2P_{n,n-3}'^{(q^2)} & \cdots & 0 \end{pmatrix} \\ \equiv & \text{Pf}(P)(\text{mod } 2), \end{aligned}$$

which implies

$$\det(B) \equiv 2S + \text{Pf}(P)(\text{mod } 4),$$

where $S \in \mathbb{Z}[X_1, \dots, X_n]$, and

$$(\det(B))^2 \equiv 4S^2 + 4\text{Pf}(P)S + (\text{Pf}(P))^2(\text{mod } 8).$$

On the other hand, by 2.2,

$$\begin{aligned} (\det(B))^2 &= \frac{1}{\det A} \det(BA^t B) \\ &= \frac{1}{\det A} \det \begin{pmatrix} 2P_{n0} & P_{n1} & \cdots & P_{n,n-1} \\ P_{n1} & 2P_{n0}^{(q)} & \cdots & P_{n,n-2}^{(q)} \\ \cdots & \cdots & \cdots & \cdots \\ P_{n,n-1} & P_{n,n-2}^{(q)} & \cdots & 2P_{n0}^{(q^{n-1})} \end{pmatrix} \\ &= \frac{1}{\det A} (4g + \det \begin{pmatrix} 0 & P_{n1} & \cdots & P_{n,n-1} \\ P_{n1} & 0 & \cdots & P_{n,n-2}^{(q)} \\ \cdots & \cdots & \cdots & \cdots \\ P_{n,n-1} & P_{n,n-2}^{(q)} & \cdots & 0 \end{pmatrix}), \end{aligned}$$

with some $g \in \mathbb{Z}[P_{n0}, P_{n0}^{(q)}, \dots, P_{n0}^{(q^{n-1})}, P]$. Thus,

$$\begin{aligned} & 4g + \det \begin{pmatrix} 0 & P_{n1} & \cdots & P_{n,n-1} \\ P_{n1} & 0 & \cdots & P_{n,n-2}^{(q)} \\ \cdots & \cdots & \cdots & \cdots \\ P_{n,n-1} & P_{n,n-2}^{(q)} & \cdots & 0 \end{pmatrix} \\ \equiv & \det(A)(4S^2 + 4\text{Pf}(P)S + (\text{Pf}(P))^2)(\text{mod } 8) \\ \equiv & ((-1)^{\nu-1}4\alpha^2 + (-1)^\nu)(4S^2 + 4\text{Pf}(P)S + (\text{Pf}(P))^2)(\text{mod } 8) \\ \equiv & (-1)^\nu(4S^2 + 4\text{Pf}(P)S + (\text{Pf}(P))^2) + (-1)^{\nu-1}4\alpha^2(\text{Pf}(P))^2(\text{mod } 8). \end{aligned}$$

But, by 2.3,

$$(\text{Pf}(P))^2 - (-1)^\nu \det \begin{pmatrix} 0 & P_{n1} & \cdots & P_{n,n-1} \\ P_{n1} & 0 & \cdots & P_{n,n-2}^{(q)} \\ \cdots & \cdots & \cdots & \cdots \\ P_{n,n-1} & P_{n,n-2}^{(q)} & \cdots & 0 \end{pmatrix} \in 4\mathbb{Z}[P].$$

Hence $S^2 + \text{Pf}(P)S \equiv f(\text{mod } 2)$, for some $f \in \mathbb{Z}[\alpha, P_{n0}, P_{n0}^{(q)}, \dots, P_{n0}^{(q^{n-1})}, P]$, as required.

□

Now we can prove the main theorem.

Theorem 3.3. *Let $\mathbb{F}_q$ be a finite field of characteristic two, $n = 2\nu$, G be $G_{2\nu}$ or $G_{2(\nu-1)+2}$, and*

$$O_n(\mathbb{F}_q, G) = \{ T \in \text{GL}_n(\mathbb{F}_q) : TG^tT \equiv G \}.$$

Let $A = G + {}^tG$ and define P_{nk} , $k = 0, 1, \dots$, as in (1) and (2). Then

$$\mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)} = \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{n, n-1}).$$

Proof. Let

$$G_A = \left\{ T \in \text{GL}_n(\mathbb{F}_q) : TA^tT = A \right\}.$$

Then $G_A \cong Sp_{2\nu}(\mathbb{F}_q)$. Let $T \in O_n(\mathbb{F}_q, G)$. Then $TG^tT + G$ is alternate, hence,

$$\begin{aligned} TA^tT + A &= T(G + {}^tG)^tT + (G + {}^tG) \\ &= {}^t(TG^tT + G) + (TG^tT + G) \\ &= 0. \end{aligned}$$

Then $T \in G_A$, thus $O_n(\mathbb{F}_q, G) \subseteq G_A$. It follows that

$$\mathbb{F}_q(X_1, \dots, X_n)^{G_A} \subseteq \mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)}.$$

It is known, as mentioned in the introduction, that $\mathbb{F}_q(X_1, \dots, X_n)^{G_A} = \mathbb{F}_q(P_{n1}, \dots, P_{nn})$. Furthermore, note that, for any $T \in \text{GL}_n(\mathbb{F}_q)$,

$$\begin{aligned} T \in O_n(\mathbb{F}_q, G) &\Leftrightarrow (X_1, \dots, X_n)(TG^tT + G) \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix} = 0 \\ &\Leftrightarrow \sigma_T(P_{n0}) = P_{n0}. \end{aligned}$$

Hence

$$\begin{aligned} \mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)} &= \mathbb{F}_q(X_1, \dots, X_n)^{G_A}(P_{n0}) \\ &= \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{nn}). \end{aligned}$$

It remains to show that $P_{nn} \in \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{n, n-1})$.

Suppose we work temporarily in $\mathbb{Z}[X_1, \dots, X_n]$ as we did before this theorem. Let

$$\tilde{P} = \begin{pmatrix} P_{n1} & P_{n2} & \dots & P_{n, n-1} & P_{nn} \\ 2P_{n0}^{(q)} & P_{n1}^{(q)} & \dots & P_{n, n-2}^{(q)} & P_{n, n-1}^{(q)} \\ P_{n1}^{(q)} & 2P_{n0}^{(q^2)} & \dots & P_{n, n-3}^{(q^2)} & P_{n, n-2}^{(q^2)} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ P_{n, n-2}^{(q)} & P_{n, n-3}^{(q^2)} & \dots & 2P_{n0}^{(q^{n-1})} & P_{n1}^{(q^{n-1})} \end{pmatrix},$$

then $BA^t B^{(q)} = \tilde{P}$. Note that

$$\begin{aligned} \det(\tilde{P}) &= \det \begin{pmatrix} P_{n1} & P_{n2} & \cdots & P_{n,n-1} & 0 \\ 2P_{n0}^{(q)} & P_{n1}^{(q)} & \cdots & P_{n,n-2}^{(q)} & P_{n,n-1}^{(q)} \\ P_{n1}^{(q)} & 2P_{n0}^{(q^2)} & \cdots & P_{n,n-3}^{(q^2)} & P_{n,n-2}^{(q^2)} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ P_{n,n-2}^{(q)} & P_{n,n-3}^{(q^2)} & \cdots & 2P_{n0}^{(q^{n-1})} & P_{n1}^{(q^{n-1})} \end{pmatrix} \\ &\quad - P_{nn} \det \begin{pmatrix} 2P_{n0}^{(q)} & P_{n1}^{(q)} & \cdots & P_{n,n-2}^{(q)} \\ P_{n1}^{(q)} & 2P_{n0}^{(q^2)} & \cdots & P_{n,n-3}^{(q^2)} \\ \cdots & \cdots & \cdots & \cdots \\ P_{n,n-2}^{(q)} & P_{n,n-3}^{(q^2)} & \cdots & 2P_{n0}^{(q^{n-1})} \end{pmatrix} \\ &= 2f + \det(P_n) - \det(P_0)P_{nn}, \end{aligned}$$

where $f \in \mathbb{Z}[P_{n0}^{(q)}, \dots, P_{n0}^{(q^{n-1})}, P_{n1}^{(q^{n-1})}, P]$ and

$$\begin{aligned} P_n &= \begin{pmatrix} P_{n1} & P_{n2} & \cdots & P_{n,n-1} & 0 \\ 0 & P_{n1}^{(q)} & \cdots & P_{n,n-2}^{(q)} & P_{n,n-1}^{(q)} \\ P_{n1}^{(q)} & 0 & \cdots & P_{n,n-3}^{(q^2)} & P_{n,n-2}^{(q^2)} \\ \cdots & \cdots & \cdots & \cdots & \cdots \\ P_{n,n-2}^{(q)} & P_{n,n-3}^{(q^2)} & \cdots & 0 & P_{n1}^{(q^{n-1})} \end{pmatrix}, \\ P_0 &= \begin{pmatrix} 2P_{n0}^{(q)} & P_{n1}^{(q)} & \cdots & P_{n,n-2}^{(q)} \\ P_{n1}^{(q)} & 2P_{n0}^{(q^2)} & \cdots & P_{n,n-3}^{(q^2)} \\ \cdots & \cdots & \cdots & \cdots \\ P_{n,n-2}^{(q)} & P_{n,n-3}^{(q^2)} & \cdots & 2P_{n0}^{(q^{n-1})} \end{pmatrix}. \end{aligned}$$

Denote the matrix obtained from P_0 by replacing all its diagonal elements by 0 and deleting its i -th row and i -th column simultaneously by $P_{(i)}$. Then $\det(P_0) \equiv 2P_0^* \pmod{4}$ where

$$P_0^* = P_{n0}^{(q)} \det(P_{(1)}) + P_{n0}^{(q^2)} \det(P_{(2)}) + \cdots + P_{n0}^{(q^{n-1})} \det(P_{(n-1)}) + l$$

with $l \in \mathbb{Z}[P]$ such that

$$2l = \det \begin{pmatrix} 0 & P_{n1}^{(q)} & \cdots & P_{n,n-2}^{(q)} \\ P_{n1}^{(q)} & 0 & \cdots & P_{n,n-3}^{(q^2)} \\ \cdots & \cdots & \cdots & \cdots \\ P_{n,n-2}^{(q)} & P_{n,n-3}^{(q^2)} & \cdots & 0 \end{pmatrix},$$

(the existence of l follows from 2.1). Clearly, $P_0^* \in \mathbb{Z}[P_{n0}^{(q)}, \dots, P_{n0}^{(q^{n-1})}, P]$. By 3.2, $\det(B) \equiv 2S + \text{Pf}(P) \pmod{4}$, hence $\det(B^{(q)}) \equiv 2S^{(q)} + \text{Pf}(P^{(q)}) \pmod{4}$. Then, from $\det(\tilde{P}) = \det(BA^t B^{(q)})$, we have that

$$\begin{aligned} &2f + \det(P_n) - 2P_0^* P_{nn} \\ &\equiv \det(A) \det(B) \det(B^{(q)}) \pmod{4} \\ &\equiv ((-1)^{\nu-1} 4\alpha^2 + (-1)^\nu) (2S + \text{Pf}(P)) (2S^{(q)} + \text{Pf}(P^{(q)})) \pmod{4} \\ &\equiv (-1)^\nu 2(S \text{Pf}(P^{(q)}) + \text{Pf}(P) S^{(q)}) + (-1)^\nu \text{Pf}(P) \text{Pf}(P^{(q)}) \pmod{4}. \end{aligned}$$

In virtue of 3.1,

$$\begin{aligned}
\text{Pf}(P) \text{Pf}(P^{(q)}) &= \det \begin{pmatrix} 0 & P_{n1} & P_{n2} & \dots & P_{n,n-1} \\ -P_{n,n-1}^{(q)} & 0 & P_{n1}^{(q)} & \dots & P_{n,n-2}^{(q)} \\ -P_{n,n-2}^{(q^2)} & -P_{n1}^{(q)} & 0 & \dots & P_{n,n-3}^{(q^2)} \\ \dots & \dots & \dots & \dots & \dots \\ -P_{n1}^{(q^{n-1})} & -P_{n,n-2}^{(q)} & -P_{n,n-3}^{(q^2)} & \dots & 0 \end{pmatrix} \\
&= \det \begin{pmatrix} P_{n1} & P_{n2} & \dots & P_{n,n-1} & 0 \\ 0 & P_{n1}^{(q)} & \dots & P_{n,n-2}^{(q)} & P_{n,n-1}^{(q)} \\ -P_{n1}^{(q)} & 0 & \dots & P_{n,n-3}^{(q^2)} & P_{n,n-2}^{(q^2)} \\ \dots & \dots & \dots & \dots & \dots \\ -P_{n,n-2}^{(q)} & -P_{n,n-3}^{(q^2)} & \dots & 0 & P_{n1}^{(q^{n-1})} \end{pmatrix}.
\end{aligned}$$

It is easy to see, by the same argument in the proof of 2.1, that

$$\begin{aligned}
&\det \begin{pmatrix} P_{n1} & P_{n2} & \dots & P_{n,n-1} & 0 \\ 0 & P_{n1}^{(q)} & \dots & P_{n,n-2}^{(q)} & P_{n,n-1}^{(q)} \\ -P_{n1}^{(q)} & 0 & \dots & P_{n,n-3}^{(q^2)} & P_{n,n-2}^{(q^2)} \\ \dots & \dots & \dots & \dots & \dots \\ -P_{n,n-2}^{(q)} & -P_{n,n-3}^{(q^2)} & \dots & 0 & P_{n1}^{(q^{n-1})} \end{pmatrix} - \det(P_n) \\
&\in \mathbb{Z}[P_{n1}^{(q^{n-1})}, P_{n2}^{(q^{n-2})}, \dots, P_{n,n-1}^{(q)}, P].
\end{aligned}$$

Hence there exists $h \in \mathbb{Z}[P_{n1}^{(q^{n-1})}, P_{n2}^{(q^{n-2})}, \dots, P_{n,n-1}^{(q)}, P]$ such that $\det(P_n) - (-1)^\nu \text{Pf}(P) \text{Pf}(P^{(q)}) = 2h$. Then

$$P_0^* P_{nn} \equiv f + (-1)^{\nu-1} (S \text{Pf}(P^{(q)}) + \text{Pf}(P) S^{(q)}) + h \pmod{2}.$$

Now pass to $\mathbb{Z}_2[X_1, \dots, X_n]$, hence, to $\mathbb{F}_q[X_1, \dots, X_n]$, thus, α is an element of $\mathbb{F}_q$. Then we have

$$\begin{aligned}
P_0^* P_{nn} &= f + h + S \text{Pf}(P)^q + \text{Pf}(P) S^q \\
&= f + h + S(S + \text{Pf}(P))^q + S^q(S + \text{Pf}(P)),
\end{aligned}$$

where $P_0^*, f, h \in \mathbb{Z}[P_{n0}, P_{n1}, \dots, P_{n,n-1}]$.

Note that, as a symmetric polynomial in S and $S + \text{Pf}(P)$, $S(S + \text{Pf}(P))^q + S^q(S + \text{Pf}(P))$ is a polynomial of $S + (S + \text{Pf}(P)) = \text{Pf}(P)$ and $S(S + \text{Pf}(P))$. But $S(S + \text{Pf}(P)) \in \mathbb{Z}_2[P_{n0}, P_{n1}, \dots, P_{n,n-1}]$ by 3.2. Hence $S(S + \text{Pf}(P))^q + S^q(S + \text{Pf}(P)) \in \mathbb{Z}_2[P_{n0}, P_{n1}, \dots, P_{n,n-1}]$.

Finally, let us show that $P_0^* \neq 0$. When $n = 2$, it is clear that $P_0^* = P_{20}^q \neq 0$. Suppose that $n > 2$. We distinguish the cases $G = G_{2\nu}$ and $G = G_{2(\nu-1)+2}$. When $G = G_{2\nu}$, set $X_\nu = X_n$ in P_0^* , then $P_{n0} = P_{n-2,0} + X_n^2$, $P_{ni} = P_{n-2,i}$, $i = 1, 2, \dots$, where $P_{n-2,i}$ is defined relative to $G_{2(\nu-1)}$ and the indeterminates $X_1, \dots, X_{\nu-1}, X_{\nu+1}, \dots, X_{2\nu-1}$. Hence $l, \det(P_{(i)}) \in \mathbb{Z}[X_1, \dots, X_{\nu-1}, X_{\nu+1}, \dots, X_{n-1}]$, $i = 1, \dots, n-1$. As a polynomial in X_n ,

the leading coefficient of P_0^* is $\det(P_{(n-1)})$. But

$$\begin{aligned}
& \det(P_{(n-1)}) \\
&= \det \begin{pmatrix} 0 & P_{n-2,1}^q & \cdots & P_{n-2,n-3}^q \\ P_{n-2,1}^q & 0 & \cdots & P_{n-2,n-4}^{q^2} \\ & \cdots & \cdots & \\ P_{n-2,n-3}^q & P_{n-2,n-4}^{q^2} & \cdots & 0 \end{pmatrix} \\
&= \det \left\{ \begin{pmatrix} X_1^q & X_2^q & \cdots & X_{n-2}^q \\ X_1^{q^2} & X_2^{q^2} & \cdots & X_{n-2}^{q^2} \\ & \cdots & \cdots & \\ X_1^{q^{n-2}} & X_2^{q^{n-2}} & \cdots & X_{n-2}^{q^{n-2}} \end{pmatrix} \begin{pmatrix} 0 & I^{(\nu-1)} \\ I^{(\nu-1)} & 0 \end{pmatrix} \begin{pmatrix} X_1^q & X_1^{q^2} & \cdots & X_1^{q^{n-2}} \\ X_2^q & X_2^{q^2} & \cdots & X_2^{q^{n-2}} \\ & \cdots & \cdots & \\ X_{n-2}^q & X_{n-2}^{q^2} & \cdots & X_{n-2}^{q^{n-2}} \end{pmatrix} \right\} \\
&= \det \begin{pmatrix} X_1 & X_2 & \cdots & X_{n-2} \\ X_1^q & X_2^q & \cdots & X_{n-2}^q \\ & \cdots & \cdots & \\ X_1^{q^{n-3}} & X_2^{q^{n-3}} & \cdots & X_{n-2}^{q^{n-3}} \end{pmatrix}^{q^2} \\
&\neq 0.
\end{aligned}$$

Hence $P_0^* \neq 0$. Then $P_{nn} \in \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{n,n-1})$. When $G = G_{2(\nu-1)+2}$, we take

$$G = \begin{pmatrix} \alpha & 1 \\ & \alpha \\ & & 0 & I^{(\nu-1)} \\ & & & 0 \end{pmatrix} \text{ and set } X_{\nu+1} = X_n, \text{ then we also have } P_{n0} = P_{n-2,0} + X_n^2,$$

$$P_{ni} = P_{n-2,i}, i = 1, 2, \dots, \text{ but now } P_{n-2,i} \text{ is defined relative to } \begin{pmatrix} \alpha & 1 \\ & \alpha \\ & & 0 & I^{(\nu-2)} \\ & & & 0 \end{pmatrix} \text{ and}$$

the indeterminates $X_1, \dots, X_\nu, X_{\nu+2}, \dots, X_{2\nu-1}$. Thus we can draw the same conclusion $P_{nn} \in \mathbb{F}_q(P_{n0}, P_{n1}, \dots, P_{n,n-1})$. $\square$

Now let us come to the case when $n = 2\nu + 1$ is odd. We need the following lemma.

Lemma 3.4. *Let $n = 2\nu$ and $\mathbb{F}_q$ a finite field of characteristic two. In $\mathbb{F}_q[X_1, \dots, X_n]$, let*

$$\begin{aligned}
P_{nk} &= (X_1, \dots, X_n) \begin{pmatrix} 0 & I^{(\nu)} \\ I^{(\nu)} & 0 \end{pmatrix} \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_n^{q^k} \end{pmatrix}, \quad k = 1, 2, \dots, \\
D_{i_1 i_2 \dots i_n} &= \det \begin{pmatrix} X_1^{q^{i_1}} & X_1^{q^{i_2}} & \cdots & X_1^{q^{i_n}} \\ X_2^{q^{i_1}} & X_2^{q^{i_2}} & \cdots & X_2^{q^{i_n}} \\ & \cdots & \cdots & \\ X_n^{q^{i_1}} & X_n^{q^{i_2}} & \cdots & X_n^{q^{i_n}} \end{pmatrix}, \quad 0 \leq i_1 < i_2 < \cdots < i_n.
\end{aligned}$$

Then

$$D_{i_1 i_2 \dots i_n} \in \mathbb{F}_q[P_{n1}, P_{n2}, \dots, P_{n,i_n-i_1}].$$

Proof. Since

$$\begin{aligned}
(D_{i_1 i_2 \dots i_n})^2 &= \det \left\{ \begin{pmatrix} X_1^{q^{i_1}} & X_2^{q^{i_1}} & \dots & X_n^{q^{i_1}} \\ X_1^{q^{i_2}} & X_2^{q^{i_2}} & \dots & X_n^{q^{i_2}} \\ \dots & \dots & \dots & \dots \\ X_1^{q^{i_n}} & X_2^{q^{i_n}} & \dots & X_n^{q^{i_n}} \end{pmatrix} \begin{pmatrix} 0 & I^{(\nu)} \\ I^{(\nu)} & 0 \end{pmatrix} \begin{pmatrix} X_1^{q^{i_1}} & X_1^{q^{i_2}} & \dots & X_1^{q^{i_n}} \\ X_2^{q^{i_1}} & X_2^{q^{i_2}} & \dots & X_2^{q^{i_n}} \\ \dots & \dots & \dots & \dots \\ X_n^{q^{i_1}} & X_n^{q^{i_2}} & \dots & X_n^{q^{i_n}} \end{pmatrix} \right\} \\
&= \det \begin{pmatrix} 0 & P_{n,i_2-i_1}^{q^{i_1}} & P_{n,i_3-i_1}^{q^{i_1}} & \dots & P_{n,i_n-i_1}^{q^{i_1}} \\ P_{n,i_2-i_1}^{q^{i_1}} & 0 & P_{n,i_3-i_2}^{q^{i_2}} & \dots & P_{n,i_n-i_2}^{q^{i_2}} \\ P_{n,i_3-i_1}^{q^{i_1}} & P_{n,i_3-i_2}^{q^{i_2}} & 0 & \dots & P_{n,i_n-i_3}^{q^{i_3}} \\ \dots & \dots & \dots & \dots & \dots \\ P_{n,i_n-i_1}^{q^{i_1}} & P_{n,i_n-i_2}^{q^{i_2}} & P_{n,i_n-i_3}^{q^{i_3}} & \dots & 0 \end{pmatrix},
\end{aligned}$$

we have that

$$D_{i_1 i_2 \dots i_n} = \text{Pf} \begin{pmatrix} 0 & P_{n,i_2-i_1}^{q^{i_1}} & P_{n,i_3-i_1}^{q^{i_1}} & \dots & P_{n,i_n-i_1}^{q^{i_1}} \\ P_{n,i_2-i_1}^{q^{i_1}} & 0 & P_{n,i_3-i_2}^{q^{i_2}} & \dots & P_{n,i_n-i_2}^{q^{i_2}} \\ P_{n,i_3-i_1}^{q^{i_1}} & P_{n,i_3-i_2}^{q^{i_2}} & 0 & \dots & P_{n,i_n-i_3}^{q^{i_3}} \\ \dots & \dots & \dots & \dots & \dots \\ P_{n,i_n-i_1}^{q^{i_1}} & P_{n,i_n-i_2}^{q^{i_2}} & P_{n,i_n-i_3}^{q^{i_3}} & \dots & 0 \end{pmatrix} \in \mathbb{F}_q[P_{n1}, P_{n2}, \dots, P_{n,i_n-i_1}].$$

□

Theorem 3.5. Let $\mathbb{F}_q$ be a finite field of characteristic two, $n = 2\nu + 1$, $G = G_{2\nu+1}$ and

$$O_n(\mathbb{F}_q, G) = \{ T \in \text{GL}_n(\mathbb{F}_q) : TG^t T \equiv G \}.$$

Let

$$\begin{aligned}
P_{n-1,0} &= (X_1, \dots, X_{n-1}) \begin{pmatrix} 0 & I^{(\nu)} \\ 0 & 0 \end{pmatrix} \begin{pmatrix} X_1 \\ \vdots \\ X_{n-1} \end{pmatrix} \\
P_{n-1,k} &= (X_1, \dots, X_{n-1}) \begin{pmatrix} 0 & I^{(\nu)} \\ I^{(\nu)} & 0 \end{pmatrix} \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_{n-1}^{q^k} \end{pmatrix}, \quad k = 1, 2, \dots.
\end{aligned}$$

Then

$$\mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)} = \mathbb{F}_q(P_{n-1,0}, P_{n-1,1}, \dots, P_{n-1,n-2}, X_n).$$

Proof. By Dickson's Theorem,

$$\mathbb{F}_q(X_1, \dots, X_n)^{\text{GL}_n(\mathbb{F}_q)} = \mathbb{F}_q(C_{n0}, C_{n1}, \dots, C_{n,n-1}),$$

where $C_{ni} = \frac{D_{ni}}{D_{nn}}$, $i = 0, 1, \dots, n-1$, and

$$D_{ni} = \det \begin{pmatrix} X_1 & X_1^q & \dots & \widehat{X_1^{q^i}} & \dots & X_1^{q^n} \\ X_2 & X_2^q & \dots & \widehat{X_2^{q^i}} & \dots & X_2^{q^n} \\ \dots & \dots & \dots & \dots & \dots & \dots \\ X_n & X_n^q & \dots & \widehat{X_n^{q^i}} & \dots & X_n^{q^n} \end{pmatrix}, \quad i = 0, 1, \dots, n.$$

Let

$$P_{n0} = (X_1, \dots, X_n)G \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix}.$$

Note that, for any $T \in \text{GL}_n(\mathbb{F}_q)$,

$$\begin{aligned} T \in O_n(\mathbb{F}_q, G) &\Leftrightarrow (X_1, \dots, X_n)(TG^tT + G) \begin{pmatrix} X_1 \\ \vdots \\ X_n \end{pmatrix} = 0 \\ &\Leftrightarrow \sigma_T(P_{n0}) = P_{n0}. \end{aligned}$$

Hence

$$\begin{aligned} \mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)} &= \mathbb{F}_q(X_1, \dots, X_n)^{\text{GL}_n(\mathbb{F}_q)}(P_{n0}) \\ &= \mathbb{F}_q(P_{n0}, C_{n0}, C_{n1}, \dots, C_{n,n-1}). \end{aligned}$$

It is easy to see that $P_{n-1,1}, \dots, P_{n-1,n} \in \mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)}$. By [8, Theorem 7.1], every $T \in O_n(\mathbb{F}_q, G)$ has the form

$$\begin{pmatrix} A & B & E \\ C & D & F \\ 0 & 0 & 1 \end{pmatrix},$$

where $\begin{pmatrix} A & B \\ C & D \end{pmatrix} \in \text{Sp}_{2\nu}(\mathbb{F}_q)$, it follows that $\sigma_T(X_n) = X_n$ which implies that $X_n \in \mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)}$. Then $P_{n-1,0} = P_{n0} + X_n^2 \in \mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)}$. In virtue of 3.4 and

$$\begin{aligned} D_{ni} &= \sum_{j=0}^{i-1} X_n^{q^j} \det \begin{pmatrix} X_1 & X_1^q & \dots & \widehat{X_1^{q^j}} & \dots & \widehat{X_1^{q^i}} & \dots & X_1^{q^n} \\ X_2 & X_2^q & \dots & \widehat{X_2^{q^j}} & \dots & \widehat{X_2^{q^i}} & \dots & X_2^{q^n} \\ & & \dots & & & & & \\ X_{n-1} & X_{n-1}^q & \dots & \widehat{X_{n-1}^{q^j}} & \dots & \widehat{X_{n-1}^{q^i}} & \dots & X_{n-1}^{q^n} \end{pmatrix} \\ &\quad + \sum_{j=i+1}^n X_n^{q^j} \det \begin{pmatrix} X_1 & X_1^q & \dots & \widehat{X_1^{q^i}} & \dots & \widehat{X_1^{q^j}} & \dots & X_1^{q^n} \\ X_2 & X_2^q & \dots & \widehat{X_2^{q^i}} & \dots & \widehat{X_2^{q^j}} & \dots & X_2^{q^n} \\ & & \dots & & & & & \\ X_{n-1} & X_{n-1}^q & \dots & \widehat{X_{n-1}^{q^i}} & \dots & \widehat{X_{n-1}^{q^j}} & \dots & X_{n-1}^{q^n} \end{pmatrix}, \\ i &= 0, 1, \dots, n. \end{aligned}$$

we see that, $D_{ni} \in \mathbb{F}_q[P_{n-1,1}, \dots, P_{n-1,n}, X_n]$, $i = 0, 1, \dots, n$. But, as seen in the introduction, $P_{n-1,n} \in \mathbb{F}_q(P_{n-1,1}, \dots, P_{n-1,n-1})$ and, from the proof of 3.3, we see that $P_{n-1,n-1} \in \mathbb{F}_q(P_{n-1,0}, P_{n-1,1}, \dots, P_{n-1,n-2})$. Hence $D_{ni} \in \mathbb{F}_q(P_{n-1,0}, P_{n-1,1}, \dots, P_{n-1,n-2}, X_n)$, hence, $C_{ni} \in \mathbb{F}_q(P_{n-1,0}, P_{n-1,1}, \dots, P_{n-1,n-2}, X_n)$, $i = 0, 1, \dots, n-1$. Therefore,

$$\mathbb{F}_q(X_1, \dots, X_n)^{O_n(\mathbb{F}_q, G)} = \mathbb{F}_q(P_{n-1,0}, P_{n-1,1}, \dots, P_{n-1,n-2}, X_n).$$

□

4. RATIONAL INVARIANTS OF PSEUDO-SYMPLECTIC GROUPS

Let $\mathbb{F}_q$ be a finite field of characteristic two. In this section, we will discuss the rational invariants of pseudo-symplectic groups.

Theorem 4.1. *Let $n = 2\nu + 1$, $S = S_{2\nu+1}$ and*

$$P_{n-1,k} = (X_1, \dots, X_{n-1}) \begin{pmatrix} 0 & I^{(\nu)} \\ I^{(\nu)} & 0 \end{pmatrix} \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_{n-1}^{q^k} \end{pmatrix}, \quad k = 1, 2, \dots$$

Then

$$\mathbb{F}_q(X_1, \dots, X_n)^{Ps_n(\mathbb{F}_q, S)} = \mathbb{F}_q(P_{n-1,1}, \dots, P_{n-1,n-1}, X_n).$$

Proof. Let

$$\overline{P}_{nk} = (X_1, \dots, X_n) S \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_n^{q^k} \end{pmatrix}, \quad k = 0, 1, 2, \dots$$

Then, as mentioned in the introduction,

$$\mathbb{F}_q(X_1, \dots, X_n)^{Ps_n(\mathbb{F}_q, S)} = \mathbb{F}_q(\overline{P}_{n0}, \overline{P}_{n1}, \dots, \overline{P}_{nn}).$$

Note that $\overline{P}_{n0} = X_n^2$ and, for any $T \in Ps_n(\mathbb{F}_q, S)$, from $\sigma_T(\overline{P}_{n0}) = \overline{P}_{n0}$, i.e., $(\sigma_T(X_n))^2 = X_n^2$, we see that $\sigma_T(X_n) = X_n$, thus $X_n \in \mathbb{F}_q(X_1, \dots, X_n)^{Ps_n(\mathbb{F}_q, S)}$. Since $\overline{P}_{nk} = P_{n-1,k} + X_n^{q^k+1}$, it follows that

$$\mathbb{F}_q(X_1, \dots, X_n)^{Ps_n(\mathbb{F}_q, S)} = \mathbb{F}_q(P_{n-1,1}, \dots, P_{n-1,n}, X_n).$$

If we set

$$A = \begin{pmatrix} 0 & I^{(\nu)} \\ I^{(\nu)} & 0 \end{pmatrix},$$

then, as mentioned in the introduction,

$$P_{n-1,n} \in \mathbb{F}_q(X_1, \dots, X_{n-1})^{G_A} = \mathbb{F}_q(P_{n-1,1}, \dots, P_{n-1,n-1}).$$

Hence $\mathbb{F}_q(X_1, \dots, X_n)^{Ps_n(\mathbb{F}_q, S)} = \mathbb{F}_q(P_{n-1,1}, \dots, P_{n-1,n-1}, X_n)$. $\square$

Lemma 4.2. *Let $f_{ij} \in \mathbb{F}_q[X_1, \dots, X_n]$, $1 \leq i \leq j \leq m$. Then there exists $f \in \mathbb{F}_q[f_{ij}, 1 \leq i \leq j \leq m]$ such that*

$$\det \begin{pmatrix} f_{11}^2 & f_{12} & \dots & f_{1m} \\ f_{12} & f_{22}^2 & \dots & f_{2m} \\ \dots & \dots & \dots & \dots \\ f_{1m} & f_{2m} & \dots & f_{mm}^2 \end{pmatrix} = f^2.$$

Proof. Let

$$B = \begin{pmatrix} f_{11}^2 & f_{12} & \dots & f_{1m} \\ f_{12} & f_{22}^2 & \dots & f_{2m} \\ \dots & \dots & \dots & \dots \\ f_{1m} & f_{2m} & \dots & f_{mm}^2 \end{pmatrix}.$$

Let r be the number of nonzero elements in the diagonal of B . We prove the lemma by induction on r . If $r = 0$, then B is alternate, hence $\det(B) = (\text{Pf}(B))^2$. Now assume that $r > 0$. We may assume that $f_{11} \neq 0$. Then $\det(B) = f_{11}^2 \det(B_1) + \det(B_2)$, where

$$B_1 = \begin{pmatrix} f_{22}^2 & f_{23} & \cdots & f_{2m} \\ f_{23} & f_{33}^2 & \cdots & f_{3m} \\ \cdots & \cdots & \cdots & \cdots \\ f_{2m} & f_{3m} & \cdots & f_{mm}^2 \end{pmatrix}, \quad B_2 = \begin{pmatrix} 0 & f_{12} & \cdots & f_{1m} \\ f_{12} & f_{22}^2 & \cdots & f_{2m} \\ \cdots & \cdots & \cdots & \cdots \\ f_{1m} & f_{2m} & \cdots & f_{mm}^2 \end{pmatrix}.$$

By induction assumption, $\det(B_1) = f_1^2$, $\det(B_2) = f_2^2$, for some $f_1, f_2 \in \mathbb{F}_q[f_{ij}, 1 \leq i \leq j \leq m]$. Let $f = f_{11}f_1 + f_2$. Then $\det(B) = f^2$, as required. $\square$

Theorem 4.3. *Let $n = 2\nu + 2$, $S = S_{2\nu+2}$ and*

$$P_{nk} = (X_1, \dots, X_n) S \begin{pmatrix} X_1^{q^k} \\ \vdots \\ X_n^{q^k} \end{pmatrix}, \quad k = 0, 1, 2, \dots.$$

Then

$$\mathbb{F}_q(X_1, \dots, X_n)^{P_{Sn}(\mathbb{F}_q, S)} = \mathbb{F}_q(P_{n1}, \dots, P_{n,n-1}, X_n).$$

Proof. As in the proof of 4.1, we have

$$\mathbb{F}_q(X_1, \dots, X_n)^{P_{Sn}(\mathbb{F}_q, S)} = \mathbb{F}_q(P_{n1}, \dots, P_{nn}, X_n).$$

Let

$$D_n = \det \begin{pmatrix} X_1 & X_2 & \cdots & X_n \\ X_1^q & X_2^q & \cdots & X_n^q \\ \cdots & \cdots & \cdots & \cdots \\ X_1^{q^{n-1}} & X_2^{q^{n-1}} & \cdots & X_n^{q^{n-1}} \end{pmatrix}.$$

Then

$$\begin{aligned} D_n^2 &= \det \left\{ \begin{pmatrix} X_1 & X_2 & \cdots & X_n \\ X_1^q & X_2^q & \cdots & X_n^q \\ \cdots & \cdots & \cdots & \cdots \\ X_1^{q^{n-1}} & X_2^{q^{n-1}} & \cdots & X_n^{q^{n-1}} \end{pmatrix} S \begin{pmatrix} X_1 & X_1^q & \cdots & X_1^{q^{n-1}} \\ X_2 & X_2^q & \cdots & X_2^{q^{n-1}} \\ \vdots & \vdots & & \vdots \\ X_n & X_n^q & \cdots & X_n^{q^{n-1}} \end{pmatrix} \right\} \\ &= \det \begin{pmatrix} P_{n0} & P_{n1} & \cdots & P_{n,n-1} \\ P_{n1} & P_{n0}^q & \cdots & P_{n,n-2}^q \\ \cdots & \cdots & \cdots & \cdots \\ P_{n,n-1} & P_{n,n-2}^q & \cdots & P_{n0}^{q^{n-1}} \end{pmatrix} \end{aligned}$$

and $P_{n0} = X_n^2$. It follows from 4.2 that $D_n^2 = f^2$ for some $f \in \mathbb{F}_q[P_{n1}, \dots, P_{n,n-1}, X_n]$. Hence $D_n \in \mathbb{F}_q[P_{n1}, \dots, P_{n,n-1}, X_n]$. On the other hand, we have

$$\begin{aligned} D_n^{q+1} &= \det \left\{ \begin{pmatrix} X_1^q & X_2^q & \dots & X_n^q \\ X_1^{q^2} & X_2^{q^2} & \dots & X_n^{q^2} \\ \vdots & \vdots & \ddots & \vdots \\ X_1^{q^n} & X_2^{q^n} & \dots & X_n^{q^n} \end{pmatrix} S \begin{pmatrix} X_1 & X_1^q & \dots & X_1^{q^{n-1}} \\ X_2 & X_2^q & \dots & X_2^{q^{n-1}} \\ \vdots & \vdots & \ddots & \vdots \\ X_n & X_n^q & \dots & X_n^{q^{n-1}} \end{pmatrix} \right\} \\ &= \det \begin{pmatrix} P_{n1} & P_{n0}^q & P_{n1}^q & \dots & P_{n,n-2}^q \\ P_{n2} & P_{n1}^q & P_{n0}^{q^2} & \dots & P_{n,n-3}^{q^2} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ P_{n,n-1} & P_{n,n-2}^q & P_{n,n-3}^{q^2} & \dots & P_{n0}^{q^{n-1}} \\ P_{nn} & P_{n,n-1}^q & P_{n,n-2}^{q^2} & \dots & P_{n1}^{q^{n-1}} \end{pmatrix}. \end{aligned}$$

Let

$$K = \det \begin{pmatrix} P_{n0} & P_{n1} & \dots & P_{n,n-2} \\ P_{n1} & P_{n0}^q & \dots & P_{n,n-3}^q \\ \vdots & \vdots & \ddots & \vdots \\ P_{n,n-2} & P_{n,n-3}^q & \dots & P_{n0}^{q^{n-2}} \end{pmatrix}.$$

Then $D_n^{q+1} = K^q P_{nn} + g$, for some $g \in \mathbb{F}_q[P_{n1}, \dots, P_{n,n-1}]$. Note that, $K \in \mathbb{F}_q[P_{n1}, \dots, P_{n,n-2}, X_n]$. Furthermore, since

$$\begin{aligned} &K|_{X_{n-1}=X_n} \\ &= \det \left\{ \begin{pmatrix} X_1 & \dots & X_{n-2} & X_{n-1} & X_{n-1} \\ X_1^q & \dots & X_{n-2}^q & X_{n-1}^q & X_{n-1}^q \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ X_1^{q^{n-2}} & \dots & X_{n-2}^{q^{n-2}} & X_{n-1}^{q^{n-2}} & X_{n-1}^{q^{n-2}} \end{pmatrix} S \begin{pmatrix} X_1 & X_1^q & \dots & X_1^{q^{n-2}} \\ \vdots & \vdots & \ddots & \vdots \\ X_{n-2} & X_{n-2}^q & \dots & X_{n-2}^{q^{n-2}} \\ X_{n-1} & X_{n-1}^q & \dots & X_{n-1}^{q^{n-2}} \\ X_{n-1} & X_{n-1}^q & \dots & X_{n-1}^{q^{n-2}} \end{pmatrix} \right\} \\ &= \det \left\{ \begin{pmatrix} X_1 & \dots & X_{n-2} & X_{n-1} & 0 \\ X_1^q & \dots & X_{n-2}^q & X_{n-1}^q & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ X_1^{q^{n-2}} & \dots & X_{n-2}^{q^{n-2}} & X_{n-1}^{q^{n-2}} & 0 \end{pmatrix} \begin{pmatrix} 0 & I^{(\nu)} & & & \\ I^{(\nu)} & 0 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} \begin{pmatrix} X_1 & X_1^q & \dots & X_1^{q^{n-2}} \\ \vdots & \vdots & \ddots & \vdots \\ X_{n-2} & X_{n-2}^q & \dots & X_{n-2}^{q^{n-2}} \\ X_{n-1} & X_{n-1}^q & \dots & X_{n-1}^{q^{n-2}} \\ X_{n-1} & X_{n-1}^q & \dots & X_{n-1}^{q^{n-2}} \end{pmatrix} \right\} \\ &= \det \left\{ \begin{pmatrix} X_1 & \dots & X_{n-2} & X_{n-1} \\ X_1^q & \dots & X_{n-2}^q & X_{n-1}^q \\ \vdots & \vdots & \vdots & \vdots \\ X_1^{q^{n-2}} & \dots & X_{n-2}^{q^{n-2}} & X_{n-1}^{q^{n-2}} \end{pmatrix} \begin{pmatrix} 0 & I^{(\nu)} & & \\ I^{(\nu)} & 0 & & \\ & & 1 & \\ & & & 1 \end{pmatrix} \begin{pmatrix} X_1 & X_1^q & \dots & X_1^{q^{n-2}} \\ \vdots & \vdots & \ddots & \vdots \\ X_{n-2} & X_{n-2}^q & \dots & X_{n-2}^{q^{n-2}} \\ X_{n-1} & X_{n-1}^q & \dots & X_{n-1}^{q^{n-2}} \end{pmatrix} \right\} \\ &= \det \begin{pmatrix} X_1 & X_2 & \dots & X_{n-1} \\ X_1^q & X_2^q & \dots & X_{n-1}^q \\ \vdots & \vdots & \ddots & \vdots \\ X_1^{q^{n-2}} & X_2^{q^{n-2}} & \dots & X_{n-1}^{q^{n-2}} \end{pmatrix}^2 \\ &\neq 0, \end{aligned}$$

we see that $K \neq 0$, thus $P_{nn} \in \mathbb{F}_q(P_{n1}, \dots, P_{n,n-1}, X_n)$. Hence

$$\mathbb{F}_q(X_1, \dots, X_n)^{Ps_n(\mathbb{F}_q, S)} = \mathbb{F}_q(P_{n1}, \dots, P_{n,n-1}, X_n).$$

□

REFERENCES

- [1] N. Bourbaki. *Éléments de Mathématique, Algèbre, Chapitre 9*, Hermann, Paris, 1959.
- [2] D. Carlisle, P. H. Kropholler. Rational invariants of certain orthogonal and unitary groups, *Bull. London Math. Soc.*, **24**(1992)57-60.
- [3] H. Chu. Orthogonal group actions on rational function fields, *Bull. Inst. Math. Acad. Sinica*, **16**(1988)115-122.
- [4] H. Chu. Supplementary note on 'rational invariants of certain orthogonal and unitary groups', *Bull. London Math. Soc.*, **29**(1997)37-42.
- [5] S. D. Cohen. Rational function invariant under an orthogonal group, *Bull. London Math. Soc.*, **22**(1990)217-221.
- [6] L. E. Dickson. A fundamental system of invariants of the general modular linear group with a solution of the form problem, *Trans. Amer. Math. Soc.*, **12**(1911)75-98.
- [7] S. M. Rajaei. Rational invariants of certain orthogonal groups over finite fields of characteristic two, *Comm. Algebra*, **28**(2000)2367-2393.
- [8] Z. Wan. *Geometry of Classical Groups over Finite Fields*, 2nd edition, Science Press, Beijing/New York, 2002.

Zhongming Tang
 Department of Mathematics
 Suzhou University
 Suzhou 215006
 P. R. China
 E-mail: zmtang@suda.edu.cn

Zhe-xian Wan
 Academy of Mathematics and System Sciences
 Chinese Academy of Science
 Beijing 100080
 P. R. China
 E-mail: wan@amss.ac.cn